

DIGESTIVE DISEASE ASSOCIATES

803-324-7607

www.digestivediseaseassociates.com

General Instructions for All Endoscopy Procedure

If you have any question, please contact our office at 803-324-7607 and ask for **Procedure Scheduling**. Also you may visit our website listed above.

Diabetics: Please follow the instructions below for your procedure:

Insulin Shots: If you take insulin shots:

The **DAY BEFORE your procedure, take your usual dose of insulin.

The **EVENING BEFORE your procedure, take ½ of your usual dose of insulin.

Make sure to check your blood sugar frequently that day. If sugar falls below 70, drink sprite or 7-UP (not diet)

If you take **Diabetes Pills**:

Take **NO diabetes pills the **Night Before or Morning of Procedure**.

MORNING OF PROEDURE:

*Take **NO** insulin shots or diabetes pills.

FOR EVERYONE THE MORNING OF THE PROCEDURE:

Only take **Heart or Blood Pressure**, medication, **Seizure** medication and/or **Reflux** medication.

When to STOP Blood Thinners:

Coumadin - 3-5 days before

Pradaxa - 2-5 days before

Eliquis - 2 days before

Effient - 7 days before

Plavix or Aggrenox - 7 days before

Brilinta- 5 days before

Xarelto - 24 hours before

Aspirin- take as normal

If you take **Iron Pills**, please **STOP 4 Days** prior to your procedure.

Please note: **FAILURE** to cancel your procedure **24 hours in advance** will result in a **\$100 No Show Fee**.