

Digestive Disease Associates

170 Amendment Ave.

Rock Hill, SC 29732

www.digestivediseaseassociates.com

Sutab Prep:

The day prior to your colonoscopy. you will need to maintain a clear liquid diet. This includes items such as water, tea, apple or white grape juice, clear sodas, clear broth, plain coffee, Jello (not red or purple). Avoid milk products. Avoid any solid food.

Starting at 4pm the day before your procedure. open one bottle of 12 tablets. Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water and drink the entire amount of remaining water over the next hour. You must drink TWO more 16 ounce containers of water over the next hour. It is important that you finish this part of the prep in two hours. Bowel movements should begin about a half hour to an hour after you finish the tablets. Feelings of bloating and nausea are common but they should subside after you start having bowel movements. If needed due to nausea, you may take a 30 minute break, then start again until all of the water is complete.

5 hours prior to your procedure. open the other bottle of 12 tablets. Fill the provided container with 16 ounces of water (up to fill line). Swallow each tablet with a sip of water and drink the entire amount of remaining water over the next hour. You must drink TWO more 16 ounce containers of water over the next hour. It is important that you finish this part of the prep within two hours. Do not drink anything else prior to procedure. ****NOTHING BY MOUTH 3 HOURS PRIOR TO PROCEDURE***FAILURE TO FOLLOW THE ABOVE INSTRUCTIONS WILL RESULT IN CANCELING THE PROCEDURE.**

Please follow these instructions for best results for your procedure. Failure to follow these instructions could result in cancellation of your procedure and rescheduling of your procedure

DIGESTIVE DISEASE ASSOCIATES

803-324-7607

www.digestivediseaseassociates.com

General Instructions for All Endoscopy Procedures

If you have any questions, please contact our office at 803-324-7607 and ask for **Procedure Scheduling**. Also, you may visit our website listed above.

You must have a driver for all testing-family member/friend/neighbor-NO TRANSPORTATION SERVICES.

Diabetics and Weight Loss Patients: Please follow the instructions below for your procedure:

****Please follow the instructions below if you take these medications**

<u>Generic name</u>	<u>Brand name</u>	<u>Instructions</u>
Dulaglutide	Trulicity	Please stop 7 days prior to procedure
Exenatide(ER)	Bydureon BCise	Please stop 7 days prior to procedure
Exenatide(IR)	Byetta	Please stop 1 day prior to procedure
Liraglutide(3mg)	Saxenda	Please stop 1 day prior to procedure
Liraglutide (1.2mg/1.8mg)	Victoza	Please stop 1 day prior to procedure
Lixisenatide	Adlyxin	Please stop 1 day prior to procedure
Semaglutide	Ozempic,Wegovy	Please stop 7 days prior to procedure
Semaglutide	Rybelsus	Please stop 1 day prior to procedure
Tirzepatide	Mounjaro, Zepbound	Please stop 7 days prior to procedure
Albiglutide	Tanzeum	Please stop 7 days prior to procedure
Galcanezumab	Emgality	Please stop 7 days prior to procedure

Please consult your PCP on how to monitor blood sugar during the interim

Diabetics

****Please hold the following Medications for 3 days prior and Morning of procedure**

Brenzavvy (bexagliflozin)

Invokana (canagliflozin)

Farxiga (dapagliflozin)

Jardiance (empagliflozin)
Steglatro (ertugliflozin)

Insulin Shots: If you take insulin shots:

The **DAY BEFORE your procedure, take your usual dose of insulin.

The **EVENING BEFORE your procedure, take ½ of your usual dose of insulin.

Make sure to check your blood sugar frequently that day. If sugar falls below 70, drink sprite or 7-UP (not diet)

If you take **Diabetes Pills:**

Take **NO diabetes pills the **Night Before or Morning of Procedure.**

MORNING OF PROEDURE:

*Take **NO** insulin shots or diabetes pills.

FOR EVERYONE THE MORNING OF THE PROCEDURE:

Only take **Heart or Blood Pressure**, medication, **Seizure** medication and/or **Reflux** medication.

When to STOP Blood Thinners: Please make sure that your prescribing provider is okay with you holding the blood thinner. If not, please notify our office immediately.

Coumadin - 5 days before

Pradaxa - 3 days before

Eliquis - 2 days before

Effient - 7 days before

Plavix or Aggrenox - 7 days before

Brilinta- 5 days before

Xarelto - 2 days before

Aspirin- take as normal

If you take **Iron Pills**, please **STOP 4 Days** prior to your procedure.

Please note: **FAILURE** to cancel your procedure **24 hours in advance** will result in a **\$150 No Show Fee.**